Pesticide

YOU,'SHOULD KNOW A B O U T

California Department of **Pesticide Regulation**

1001 I Street P.O. Box 4015 Sacramento, CA 95812-4015 916-445-4300 www.cdpr.ca.gov

BRANCHES

Enforcement 916-324-4100

Northern Regional Office 916-376-8960

Central Regional Office 559-297-3511

Southern Regional Office 714-279-7690

Environmental Monitoring 916-324-4039

> **Fiscal Operations** 916-324-1350

Human Health Assessment 916-445-4233

Information Technology 916-445-2992

> Personnel 916-322-4553

Pest Management and Licensing 916-445-3914

Licensing and Certification Office

916-445-4038

Pesticide Registration 916-445-4400

Worker Health and Safety 916-445-4222

Emergency! What to do when accidents happen

Despite what many of us might think, no substance is inherently safe or unsafe. Any substance - even the most innocuous - can be harmful if you are exposed to too much of it. Pesticides are designed to be toxic to the target pest while being safe to the people that use these chemicals. However, accidents and misuse occur and they can cause illness or injury. The potential for a pesticide to cause injury depends upon several factors:

Toxicity

Toxicity is the potential a chemical, such as a pesticide, has for causing harm. Some pesticides have low human toxicity while

others are extremely toxic.

Dose

The greater the quantity of a chemical vou are exposed to, the


greater the risk of injury. The effect of a given dose varies with the weight and age of the person. Therefore, an amount that is relatively harmless to an adult may hurt a small child.

· Route of absorption

The ways the body comes in contact with chemicals. Some common routes of exposure are dermal (skin), oral (by mouth), and inhalation (breathing). Swallowing a pesticide usually creates the most serious problem. In practice, however, the most common route of absorption of pesticides is through the skin.

Duration of exposure

The longer a person is exposed, the more chemical their body absorbs.



After you have administered first aid, you should get medical help immediately. If someone develops symptoms after using or being around pesticides, call toll-free 1-800-222-1222. This will connect you to the Poison Control Center, for free medical information.



Single copies of this handout are available by calling your County Agricultural Commissioner's office, from DPR at 916-445-3974, or can be downloaded from DPR's Web site, www.cdpr.ca.gov, "Consumer Fact Sheets."

· Physical and chemical properties

Some pesticides evaporate more readily than others, so they can be more easily inhaled. Some break down quickly on surfaces, others last longer. These qualities affect the potential risk of overexposure.

· Population at risk

Persons who run the greatest danger of pesticide illness are those whose exposure is highest, such as workers who mix or apply pesticides. However, consumers who use pesticides in their homes may also be overexposed, especially if they do not follow carefully the instructions on the container label.

Recognizing pesticide poisoning

Like other chemicals, pesticides may produce injury externally or internally.

Pesticides can cause contact-associated skin irritation or allergies. Symptoms of irritation include redness, itching, or pimples. Allergic skin reactions may produce redness, swelling, or blistering. The mucous membranes of the eyes, nose, mouth, and throat are also quite sensitive to chemicals. Stinging and swelling can occur.

Internal injuries may occur depending upon where a chemical is transported in the body or what organ is affected. Shortness of breath, excessive saliva and rapid breathing may occur because of lung injury. Other symptoms to watch for include nausea, vomiting, diarrhea, headache, or dizziness.

First aid for pesticide poisoning

Appropriate first aid treatment depends on which pesticides were used. Here are some tips that may precede but should not substitute for medical advice or treatment:

Poison on skin

Irrigate the area with running water for 10 to 15 minutes. Meantime, call a physician or a poison control center for further treatment advice. Later, discard contaminated clothing or wash it thoroughly, separate from your other laundry.

· Poison in eye

Open eyelid and wash the eye slowly and gently with water. Continue eye irrigation for 10 to 15 minutes. Call a physician or a Poison Control Center for further advice.

Inhaled poison

Take the victim to fresh air. If unconscious, give artificial respiration and call for paramedic assistance. Call a physician or a poison control center for further advice.

Swallowed poison

If the person is alert and able to swallow, dilute the ingested substance with sips of milk or water. Call a physician or a Poison Control Center for what to do next.

After you have administered first aid, you should get medical help immediately. If someone develops symptoms after using or being around pesticides, call your local Poison Control Center. Dialing toll-free 1-800-222-1222, from anywhere in California, will get you free information on what to do.

Ask if the symptoms are pesticide-related and what you should do – being too cautious is better than not. Be sure to have the pesticide container when you call. Medical personnel will ask what the chemical is. Overexposure to certain chemicals causes characteristic symptoms and the doctor needs to know what the chemical is before prescribing treatment. If you are advised to seek treatment at an emergency room or physician's office, you should bring the product label to show your doctor.

To avoid problems, you should minimize your exposure when mixing and applying pesticides by wearing gloves and other protective clothing. Also be careful to follow the label instructions for mixing and application. Safe use depends on that.

Keep information handy

By calling 1-800-222-1222 from anywhere in California, you can reach a Poison Control Center.

