

AGRICULTURAL SOURCES OF DDT RESIDUES
IN CALIFORNIA'S ENVIRONMENT

September 1985

reprinted July 1996

A Report Prepared in Response
to House Resolution No. 53 (1984)

ENVIRONMENTAL HAZARDS ASSESSMENT PROGRAM

California Department of Food and Agriculture
1220 N Street
Sacramento, California 95814

AGRICULTURAL SOURCES OF
DDT RESIDUES IN
CALIFORNIA'S ENVIRONMENT

SEPTEMBER 1985

reprinted July 1996

A REPORT PREPARED IN RESPONSE
To HOUSE RESOLUTION No. 53 (1984)

BY

TOM MISCHKE
KATHY BRUNETTI
VIC ACOSTA
DON WEAVER
MARY BROWN

ENVIRONMENTAL HAZARDS ASSESSMENT PROGRAM
CALIFORNIA DEPARTMENT OF FOOD AND AGRICULTURE

1220 N STREET

SACRAMENTO, CALIFORNIA 95814

ACKNOWLEDGMENTS

The authors wish to thank the following persons for their assistance in preparing the report.

Phil Anderson	Kay Mowery
Charles Andrews	Sherman Nash
Paula Bonilla-Valdez	Liz Onstott
George Brady	Ronald Oshima
Ana Castro	Bob Rollins
Chris Collison	Lisa Ross
Dave Conrad	Nirmal Saini
James Echelberry	Roger Sava
Cindy Garretson	Scott Simpson
Linda Heath	Peter Stoddard
Maria Martinez	Roberto Teso
Nancy Miller	John Troiano
	Rob Ward

We are grateful to Richard Nutter, County Agricultural Commissioner, and Susan Cohen, Senior Deputy Agricultural Commissioner, Monterey County; Bruce Agee, State Water Resources Control Board; Paul Jaggard, Central Valley Water Quality Control Board, for their help with regard to the Blanco Drain Study in Monterey County. We are also grateful to Brian Finlayson and Norm Morgan of the California Department of Fish and Game for quality assurance work on the chemical analysis.

We would also like to thank the County Agricultural Commissioners of the following counties for their contributions to the 1985 Soil Monitoring Survey for DDT: Alameda, Colusa, Contra Costa, Fresno, Glenn, Imperial, Kern, Kings, Los Angeles, Madera, Merced, Monterey, Napa, Orange, Placer, Riverside, Sacramento, San Benito, San Bernardino, San Diego, San Joaquin, San Luis Obispo, Santa Barbara, Santa Cruz, Solano, Sonoma, Stanislaus, Sutter, Tehama, Tulare, Ventura, Yolo.

TABLE OF CONTENTS

Table of Contents i

List of Tables ii

List of Figures iii

Foreword iv

Executive Summary 1

Introduction 4

I. Agriculturally Related Sources of DDT 5

 Background and Statement of the Problem 7

 Other Pesticides as a Source of DDT 8

 Possible Illegal Use of DDT and DDD 9

 Likely Sources of DDTr Contamination from Agricultural Sources 17

II. 1985 California Soil Monitoring Survey for DDTr

 Objective and Study Plan 22

 Results 27

III. Conclusions 34

References 36

Appendix I 37

LIST OF TABLES

	Pages
Table I. DDT Use in California from 1970 to 1980	8-9
Table II. DDT and DDE Concentrations in Produce Samples Collected from 1981-84	16-21
Table III. Data from 1985 Soil Monitoring Survey for DDTr	32-35

LIST OF FIGURES

	Page
Figure 1.	Structure of Various DDTr Species 7
Figures 2-4.	Photographs of Blanco Drain, Salinas, California 26
Figures 5-7.	Photographs of Blanco Drain, Salinas, California 27
Figure 8.	Generalized Map of the Sampling Locations 30
Figure 9.	Percentage of o,p' and p,p' DDT in Total DDTr Residues for All Counties Sampled 36
Figure 10.	Percentage of o,p' DDT in Sum of o,p' DDT and p,p' DDT for All Counties Sampled 38

AGRICULTURAL SOURCES OF DDT RESIDUES IN CALIFORNIA'S ENVIRONMENT

July, 1996 Reprint

Foreword

The pesticide DDT has not been registered for use in the U.S. or California since its 1972 cancellation. This report was prepared in 1985, in response to a resolution passed by the California Legislature which directed the California Department of Food and Agriculture to investigate possible sources of DDT in the environment. (The pesticide programs of the Department of Food and Agriculture were transferred to the newly formed Department of Pesticide Regulation [DPR] in 1991.) This reprint corrects some errors previously included in an errata sheet. The report is also available at DPR's Internet Web Site at <http://cdpr.ca.gov> or you may send E-mail to info@empm.cdpr.ca.gov for further information.

Are there records showing how much DDT was applied to a specific location in California.?

Until 1971, use of DDT was not required to be reported to the California Department of Food and Agriculture, therefore, no detailed records of use exist.

What hazards may be posed by DDT residues in soil? What laws and regulations apply to areas that may be contaminated with DDT?

The California Department of Pesticide Regulation does not regulate cleanup or abatement of sites that may be contaminated by DDT as a result of previous legal use or manufacture, nor does DPR set or enforce levels for DDT in water or soil.

For additional information about current laws and regulations concerning DDT residues in soil in California, contact
California Department of Toxic Substances Control
Office of Scientific Affairs
400 P Street
Sacramento, California 95814
(916) 327-2500

For information about DDT in drinking water in California, contact
California Department of Health Services,
Division of Drinking Water, and Environmental Management
601 North 7th Street
Sacramento, California 95814
(916) 322-2308

Outside of California, contact your local agricultural, health, or environmental protection agency or the U. S. Environmental Protection Agency. You will find them listed in the government section of the white pages of your local telephone directory.

EXECUTIVE SUMMARY

HR 53, adopted by the Assembly on August 31, 1984, directed the Department of Food and Agriculture (CDFA) to investigate possible sources of DDT in the environment and to report findings to the Legislature within one year. This resolution was introduced in response to studies showing that, although its use was banned in 1973, DDT residues are still being found in California water, fish, shellfish, and produce samples. Additionally, the chemical composition of the DDT being found indicated that it might be from recent use. This report discusses CDFA's findings.

CDFA investigated three possible sources of contamination by DDT and/or its breakdown products (the combination of DDT and its breakdown products is called DDT_r) which are agriculturally related:

1. New illegal use of DDT
2. Use of other pesticides that might be contaminated with DDT_r
3. Long-lived residues from previous legal applications of DDT

The following data were analyzed in this investigation:

Field Studies

1. 1985 study, in response to HR 53, monitoring current DDT_r levels in soil from agricultural areas where DDT has historically been used.
2. 1984 case study of DDT residue levels in fish and mussels in the Salinas River, Monterey County.

Ongoing Regulatory Compliance Monitoring

- product quality monitoring of pesticides for sale, including newly introduced pesticides and new uses of existing pesticides.
- monitoring for pesticide residues in fresh produce, including fruits, nuts, vegetables, eggs, honey, hay and fodder.
- monitoring of activities of all persons selling, applying, storing, or otherwise handling pesticides.

- site inspections relating to applications, mixing and loading, equipment, storage facilities, and field worker safety.

Based on all available evidence, CDFA concluded that long-lived residues from previous applications are the apparent source of DDT residues in produce and in the environment.

Specific findings of the study include:

1. Before its ban, DDT was widely used in California in agriculture and for control of mosquitoes and other disease-carrying insects.
2. There is no evidence that there has been any illegal use of DDT since its ban. For example, in 1983, 87,000 pesticide use enforcement inspections and 3,501 investigations of possible violations were made by California County Agricultural Commissioners. None of these involved DDT. Also, in 1983, about 1300 pesticide samples were analyzed to determine what chemicals they actually contained. The results show 97.5% of these samples met registration and labeling requirements. The remaining 2.5% did not involve DDT. Even before its ban, agricultural use of DDT was declining as more insects became resistant to DDT.
3. Contamination of other pesticides by DDT can not account for the residues. There have been reports that dicofol (Kelthane[®]) contained large amounts of DDT. Samples of dicofol sold in California examined in 1983-84 contained very low levels of DDT, usually less than 1%, too low to account for DDT residues found.
4. Detectable levels of DDT are still being found on some California produce. These levels are, in most cases, well below acceptable levels. Nearly all produce samples found with residues of DDT residues have an edible portion which grows in or close to the ground, such as carrots, beets, lettuce, or spinach. DDT residues found on produce are probably the result of contamination from soil containing DDT.
5. On the average, about half the DDT detected is present as DDT in the environment. However, the composition of DDT found in soil is more stable than previously thought, therefore the kinds of DDT

residues present in soil do not necessarily indicate new use. DDT_r residues may survive in California soil for 12-15 years or more.

6. Soil contaminated with DDT_r may be moved into drains as a result of normal field work such as land leveling. Fish and shellfish pick up DDT_r from the soil particles in the water.
7. DDT_r residues are present in soil wherever DDT was used legally in the past. In 1985, CDFA collected 99 soil samples in 32 California counties from locations where DDT had been used in the past. All samples contained DDT_r.


Detectable levels of DDT_r are widespread in California soils 12 years after the ban of DDT. Based on analysis of historical and empirical evidence, CDFA concluded that residues from legal applications of DDT, before its use was banned, appear to be the source of this contamination.

I. AGRICULTURALLY RELATED SOURCES OF DDT


Background

DDT was first synthesized by Zeidler in 1874². In 1939, Dr. P. Muller working for the J.R. Geigy Chemical Company discovered the insecticidal properties of DDT³. Muller received the Nobel Prize in 1948 for this work. DDT's first full-scale uses were as the main agent for the control of insect vectors of an impressive list of diseases, including malaria, Chagas' disease, plague, typhus, yellow fever, dengue/haemorrhagic fever, encephalitis, filariasis, African trypanosomiasis, onchocerciasis and leishmaniasis⁴. In fact, even today DDT is still the main agent for control of these diseases in developing countries. Starting in 1944, DDT found widespread acceptance as a nearly universal insecticide for agricultural as well as residential, commercial and public health applications. In the late 1960's, DDT use in the United States and California reached its peak. Increasing problems with DDT resistance from a large number of insects, the development of more specific pesticides and the rising concern for environmental contamination caused a marked drop in DDT use in the United States generally, and in California in particular. In August, 1971, the United States Environmental Protection Agency (EPA) instituted official hearings on the problems caused by the continued use of DDT. These hearings continued until February 1972. The results of these hearings were given legislative backing with the elimination of use of DDT for all domestic purposes as of December 31, 1973.


The history of DDT usage in California began around 1944. DDT found wide acceptance in California as it did in the rest of the United States and the world. The uses of DDT in California ranged from control of agricultural pests to control of cockroaches in residences and mosquito abatement in neighborhoods. DDT was declared a restricted material by the California Department of Food and Agriculture in 1963. The last year in which substantial amounts of DDT were applied to California crops was 1970. Table I gives the use of DDT and DDD from 1970 to 1980. DDD is included because it too was registered for use as a pesticide.




p,p' DDT




o,p' DDT




p,p' DDE



o,p' DDE



p,p' DDD



o,p' DDD

Figure 1

Structure of Various DDT_r Species

Table L- DDT Use in California from 1970 to 1980^a

<u>Year</u>	<u>Pounds Used</u>	<u>Main Use</u>
1970	1,164,699	agricultural
1971	111,058	agricultural
1972	80,800	agricultural
1973 ^b	NUR ^{b,c}	-
1974	160	residential pest control (SLN)
1975-1980	less than 200 lbs per year	Vector control (SLN)

DDD Use in California From 1970-1980

<u>Year</u>	<u>Pounds Used</u>	<u>Main Use</u>
1970	7,929	agricultural
1971	5,601	agricultural
1972	NUR ^c	-
1973 ^b	NUR	-
1974	NUR	-
1975-1980	NUR	-

- a. 1970 was the first year in which the amount of restricted pesticides used in California was reported. In 1980, the introduction of new pesticides replaced the need to use DDT for vector control.
- b. Year all use banned except for special local needs (SLN)
- c. NUR - no use reported

Statement of the Problem

The current renewed interest in environmental DDT_r levels arising from agricultural sources is the result of continued monitoring of California's environment and agricultural products. Ongoing studies by the State Water Resources Control Board (SWRCB)⁵ have found DDT and its metabolites in fish samples taken from selected rivers in California.

The levels of DDT found in these samples were higher than expected by the SWRCB for a compound whose last major usage was fifteen years ago. Additionally, the SWRCB felt the isomeric composition of the DDT found in these samples may have been due to newly mobilized DDT residues. The SWRCB, in cooperation with the Department of Food and Agriculture conducted an intensive study of the possible sources of DDT in one of these rivers, the Salinas, in Monterey County. The full report and results of this study will be issued by the SWRCB in late 1985, but findings are briefly discussed later in this report.

The Pesticide Residue and Monitoring Program conducted by CDFA provides for the continuing inspection and sampling of farm commodities including fruits, nuts, vegetables, eggs, honey, hay and fodder. This program continues to find very low levels of DDT on some California grown produce. However, these levels of DDT are, in most cases, well below established, acceptable levels for the respective commodity. The CDFA policy on detectable levels of pesticides on commodities for which no tolerance for that pesticide has been established, or on which an over-tolerance level has been found, is to review and investigate each find on a case by case basis.

The main question that arises from the data on DDT levels in produce is the source of these DDT residues. The concentrations have remained low but are still being found some twelve years after use of the DDT compounds was banned. Possible sources considered for these DDT residues in the environment are: previous legal use; new, illegal use; or the use of other DDT containing pesticides. This report will present data on the agricultural sources of DDT in the California environment.

Other Pesticides as a Source of DDT

The one pesticide of major interest as a source of DDT residues is dicofol. Dicofol is marketed under the trade name Kelthane®. Dicofol is used to control mite pests on various commodities grown throughout the state. In order to understand the concern surrounding the use of dicofol, one must first look at the manufacturing process for dicofol. The starting material from which dicofol is made is DDE. The DDE used to make dicofol will contain some DDT as a result of its own manufacturing process. The DDE is then reacted with chlorine to form a product one can describe as "chlorinated DDE." This "chlorinated DDE" is then reacted further to give dicofol.

The final product, dicofol, therefore can contain levels of DDT and "chlorinated DDE"^{6,7}. Dicofol itself does not breakdown in the environment to DDT. However, there is evidence that the "chlorinated DDE" species will dechlorinate back to DDE in the environment.⁸ This means that dicofol can be a direct source of both DDT and DDE in the environment.

CDFA, together with the California Department of Fish and Game (CDFG) has analyzed samples of technical (undiluted) and formulated (diluted) dicofol and Kelthane[®] for DDT. CDFA drew a total of 27 samples in 1982 and 1983. Twenty-one were from registrants' formulated products and six were EPA samples of 98% technical material. Analyses revealed that one of the 21 registrants' formulated products contained under 0.1% DDT, and one of the six technical material samples contained 0.27% DDT. All the other samples contained no detectable DDT at a detection level of 0.08%. The CDFA and CDFG conducted a joint analysis of a sample of technical product in 1984 and both agencies found less than 0.1% DDT in the sample.

The manufacturer of Kelthane[®], Rohm and Haas, has announced that the current levels of DDT related impurities in Kelthane[®], will be reduced to a total of 2.5% by 1986 and to 0.1% by 1987.⁹ This reduction will mean the sum of DDT, DDE, DDD and "chlorinated DDE" will not exceed 0.1% after 1987. A level of 0.1% for the DDT related impurities in Kelthane[®] will significantly reduce the contribution of Kelthane[®] to DDT levels in the environment.

Possible Illegal Use Of DDT And DDD

California's regulatory program of pesticide use enforcement sampling, inspection, and surveillance is administered by the Pesticide Enforcement Branch of the CDFA and the County Agricultural Commissioners. This regulatory program provides lines of defense and safeguards against illegal pesticide use and residues which may occur on agricultural commodities or in the environment. These safeguards consist of field activities involving pesticide product quality sampling and inspections, pesticide residue sampling and monitoring, and pest control site inspections, as well as pest control record reviews by the County Agricultural Commissioners.

The following is an overview of field enforcement activities:

1. CDFA Pesticide Product Quality Program

The program's goal is to ensure that pesticide products offered for sale in California meet label guarantees, are registered for use in California and are unadulterated. Unregistered, adulterated or misbranded products may be quarantined from sale or use, and parties responsible for the violation may be cited or prosecuted. None of the pesticides found out of compliance in 1983 contained DDT or DDD.

The program's activities and workload consist of taking about 1,300 pesticide samples for analysis annually, primarily at dealer and user locations, and conducting about 4,000 pesticide inspections annually. Samples taken in the field are submitted to the CDFA laboratories for analysis.

"Inspection call" activities include: review of labels for compliance with labeling requirements; premise and product inspection to ensure that products are handled and packaged safely in undamaged containers; verification that products offered for sale have both current California and EPA registration; and verification that pesticide handlers hold the required licenses.

In addition, producers' establishment inspections, import inspections, and experimental use inspections are conducted by the CDFA under an enforcement agreement/grant from the U.S. Environmental Protection Agency (EPA). The breakdown of the workload and measure of effectiveness for the calendar year 1983 are shown below:

1983 WORKLOAD

<u>Type of Activity</u>	<u>Number of Samples Collected</u>
Market Surveillance	1,101
Producer Establishments	247
Inspection Calls	
Market Surveillance	3,975
Producer Establishments	194
Experimental Use Inspections	10
Import Samples & Inspections	3
Quarantine Actions	106
Notice of Warnings	66
Unregistered Product Reports	39

1983 MEASURE OF EFFECTIVENESS

Percentage of pesticides found in field inspections that are properly registered and labeled	97.5%
Percentage of pesticide formulations found in field inspections that meet the label guarantee	95.2%
Percentage of pesticide labels found in field inspections with labels in compliance with labels on file with the Department	99.2%

2. CDFA Pesticide Residue and Monitoring Program

The primary goal of this program is to protect the public and the California environment from possible harmful effects of pesticides. This goal is accomplished by two methods. The first is a produce sampling program to assure the consumer that California produce is within the legal pesticide residue tolerances established by EPA for health

and safety. The second method of accomplishing this goal is through the monitoring of newly introduced pesticides, as well as new uses of existing pesticides and special local need uses of canceled pesticides to determine that they are not creating a residue problem or having a detrimental effect on public health, wildlife, crops or the environment.

Produce inspections are primarily carried out at large wholesale markets and chain store receiving docks.

Roadside stands and retail stores are inspected through periodic surveys that are conducted in a given area. Hay and fodder are usually sampled in the areas of production. Analyses of the samples are performed by CDFA laboratories located in Sacramento, Berkeley, Fresno, and Anaheim. For special problems in growing, packing and marketing areas, a mobile laboratory is available for conducting numerous residue tests rapidly in the field.

All the pesticide residues found on these samples are entered into a data processing program that keeps a continuous record of all pesticides found in all crops. With this information, CDFA can monitor the total pesticide residue found on produce, the average level of pesticides found on produce, whether the levels are going up or down year by year, what the average percentage of residue compared to tolerance is for each pesticide and crop, and could calculate what the average intake in a person's diet would be based on individual food intake. These data may also trigger the need for reevaluation of pesticide residue data that was presented at the time the pesticide was registered for use in California.

Table II gives a summary of information on the number of produce samples with detectable DDT and DDE residues from 1981 to 1984. No DDD was detected in any samples. If illegal usage of DDT on crops were occurring in California, the Residue Program would have detected increased residues on the target crops. This has not been the case. Residue levels have remained low and have been confined to crops which either have edible portions which grow in the soil, have fruit which rests on the soil, or have an irregular leaf or fruit surface which readily traps soil particles.

Commodity	# w/ residue	Residue Found(ppm) ^b	Tolerance (ppm) ^c
1983 DDE			
Carrot	4	0.06-0.12	3.5
Endive	1	0.05	1.0
Kale	2	0.10	1.0
Lettuce (head)	1	0.02	7.0
Lettuce (leaf)	3	0.04-0.20	7.0
Mustard	1	0.05	1.0
Green Onions	1	0.11	NTE
Orange	1	0.06	3.5
Parsley	1	0.07	NTE
Parsnip	1	0.04	1.0
Radish	8	0.04-0.15	1.0
Spinach	11	0.04-0.10	1.0
Turnip	1	0.04	1.0
Zucchini	1	0.02	0.5
Sweet potatoes	1	0.03	1.0
Cilantro	1	0.10	NTE
1984 - Total of 7243 produce samples analyzed^a			
1984 DDT			
Celeriac	1	0.08	NTE
Radish	1	0.07	1.0
Spinach	2	0.14-0.18	1.0
Tomatillos	1	0.16	NTE
1984 DDE			
Beets	4	0.03-0.14	1.0
Carrots	18	0.01-0.10	3.5
Celeriac	1	0.05	NTE
Cilantro	2	0.04-0.10	NTE
Escarole	2	0.01-0.05	1.0
Lettuce (leaf)	3	0.02-0.08	7.0
Lettuce (Romaine)	2	0.03-0.18	7.0
Mustard	1	0.06	1.0
Parsley	2	0.05	NTE
Radish	15	0.02-0.15	1.0
Salsify	1	0.06	NTE
Spinach	13	0.02-0.07	1.0
Squash (summer)	2	0.03	0.50
Turnip	1	0.03	1.0

- a. No DDD detected on any samples.
b. For more than one sample analyzed, the range of results is listed.
c. Tolerances are established for DDT, DDD, and DDE either separately or as the sum of all three for each year listed.
d. NTE - No tolerance established.

3. County Agricultural Commissioner Pesticide Enforcement Program

Each county in California has a Department of Agriculture managed by a County Agricultural Commissioner. The Commissioners enforce California laws and regulations pertaining to pest control and pesticides. The commissioners are responsible for local administration of the enforcement program. CDFA is responsible for overall statewide enforcement through the issuance of policy and procedures, and by providing assistance in training, laboratory services, investigations of pesticide product quality and pesticide residues, and by providing program uniformity and coordination.

The County Agricultural Commissioners' Pesticide Enforcement Program includes: county registration of Pest Control Operators (PCO), Pest Control Advisers (PCA), aircraft pilots, and structural pest control operators (SPCO); issuance of restricted material use permits; record inspections relating to employer headquarters and PCO, pesticide dealers, PCA, and certified applicator certification records; and site inspections relating to application, mixing and loading, application equipment, storage facilities, and field worker safety; and episode investigations relating to human, environment, crop and property damage, and employee and general public complaints.

During fiscal year 1983-84, County Agricultural Commissioners expended a total of 375,274 enforcement hours. Over 87,000 pesticide use enforcement inspections were made to check compliance, and a total of 3,501 episode investigations were conducted. Administrative and judicial actions taken by the County Agricultural Commissioners were: 1470 notice of violations, 48 cease and desist orders, 72 permit actions, 3 actions against private applicator certifications, 13 county registration actions, notice to appear and 18 criminal complaints. None of these actions involved the use of DDT pesticides.

4. Pesticide Episode Reporting and Investigative Program Via Cooperative Agreements

Vital to the effectiveness of the CDFA and the County Agricultural Commissioners pesticide enforcement programs is the establishment of cooperative agreements between other county, state, and federal agencies which

The above conclusions were also reached in a SWRCB funded study of the Blanco Drain area conducted by the Bodega Marine Laboratory of the University of California.¹⁰ This study was independent of the SWRCB, CDFA, CCRWQCB, and Monterey County Agricultural Commissioners joint study.

The photographs in Figures 2-7 were taken at the Blanco Drain in late 1984. All the photographs were taken on the same day at various locations along the Blanco Drain. Figures 2 to 7 show that during post-harvest operations most fields adjacent to the drain are plowed, leveled and furrowed right up to the edge of the drain or its tributaries. After cultivation, road ways are re-established through the freshly worked ground next to the drain. During the cultivation process, excess soil is allowed to fall directly into the drain or its tributaries. Subsequent flooding of the drain or its tributaries carries this DDT_r-laden soil down the Blanco Drain to the Salinas River where it can enter the food chain of fish, mussels and other animals. This process does not occur at all points along the drain, however. Figure 7 shows an example of a grower who follows best soil management practices. The old, established roadway along the tributary was undisturbed during the cultivation operation which prevented direct movement of excess soil into the tributary. The Monterey County Agricultural Commissioner is currently informing growers in the Blanco Drain area of the results of these studies and encouraging growers to prevent soil from spilling into drains during cultivation.

This is a case study of only one of the rivers in California in which the SWRCB has found DDT_r levels. How this case study and new data collected by the CDFA relates to other rivers in California will be discussed in the conclusion section of this report.




Figure 2




Figure 3




Figure 4



Figure 5



Figure 6



Figure 7

Agricultural Produce Contamination

The Blanco Drain case study which was just discussed above serves as an excellent example for the discussion of DDT_r levels in California produce. The study found that the average DDT_r level in soils adjacent to the Blanco Drain is 2.6 ppm. CDFA collected samples of crops from fields in the Blanco Drain area in April and July, 1985. The crops sampled included cauliflower, broccoli, kale, leaf lettuce, green onions, Napa cabbage, Romaine lettuce, celery, Bok Choy, and head lettuce. No residues of DDT, DDE, or DDD were found on any of the samples at a detection limit of 0.02 to 0.03 ppm. Apparently not all produce grown in soil where low DDT_r levels exist will have residues of DDT_r. What about the produce in Table II which did show detectable levels of DDT_r? If the DDT_r found in the analysis were from DDT_r incorporated into the commodity during growth, one would expect that residues of DDT_r would always be found in every sample which had been grown in DDT_r-laden soil, and that residues would be at much higher levels than actually observed. The fact that produce grown in the DDT_r-laden soils of the Blanco Drain do not always have DDT_r residues, and that when these residues occur, they are at very low concentrations, implies that the source of DDT_r residues must be located somewhere other than in the flesh of the commodity. As discussed earlier, DDT_r residues have been found on crops which either have edible parts growing in the ground, have fruit which rests on the ground, or have an irregular leaf, stem or stalk structure which will trap soil particles. The commodities listed in Table II all fall into this category. The guidelines for analysis of these products published by the U.S. EPA under Title 40, Code of Regulations, requires that soil particles which would normally be present on the commodity when it is sold not be removed before analysis. It is possible that very small amounts of DDT_r-containing soil trapped on the commodity, in the curly leaves of parsley for example, would give a detectable level on analysis but would probably be removed by the consumer during washing prior to consumption. Further investigative work is needed to confirm this hypothesis.

II. 1985 CALIFORNIA SOIL MONITORING SURVEY FOR DDT_r

The specific objective of this monitoring study was to collect soil samples from areas in California's agricultural basins, where historic use of DDT was confirmed, in order to establish the range of ratios of DDT to total DDT_r and the range of ratios of o,p' DDT to total DDT. The Agricultural Commissioner in each county located in the

agricultural basins was contacted and with the help of information provided by his or her staff members, the degree of known use of DDT in a county was determined. If little or no DDT had been used, or the Agricultural Commissioner could not supply the needed information, soil in the county was not sampled.

For counties that had a history of moderate to high DDT use, the Agricultural Commissioner's staff was asked to provide maps or coordinates that would locate particular fields or general areas where the applications were known to have been made. This information was later used to locate sampling sites.

After the information for all selected counties had been reviewed, a decision was made to collect either two or four soil samples in a county based on the intensity of agriculture, the probable historical use of DDT, and the total land area. When more sites than required were identified in a county, an attempt was made to select sites that were most widely spaced within the county.

Soil samples were collected from individual fields with a known history of DDT application or from fields located in areas known to have had historic widespread and repeated applications of DDT. The fields selected for sampling were all under cultivation although many did not have a crop present at the time sampling was conducted. A total of 99 samples in 32 counties were collected. A generalized map of sampling locations throughout the state appears in Figure 8.

Sampling Plan

For smaller fields (about 10-30 acres), the person collecting the samples walked approximately 50 feet in from the edge, moved in a line diagonally across the field, and collected ten 1-pint soil samples at intervals along the line. Larger fields were sampled at ten points around the perimeter of the field with each sample collected about 50 feet in from the edge. A precleaned trowel was used to remove the top 6 inches of soil from a small area and place it into a clean 1-pint jar. Whenever possible, the soil was taken from a raised planting bed. Soil from the ten jars was combined in a clean three gallon stainless steel pail and thoroughly mixed using the trowel. Two 1-quart jars were then filled with soil, capped with aluminum foil-lined lids and stored at 4° C until analyzed. Each sample was accompanied by a chain of custody form which documented the county, location and date of collection. All samples were collected during April and May, 1985. One quart jar from each sampling site was analyzed by the CDFA laboratory in Sacramento for o,p' DDT, p,p' DDT, o,p' DDD, p,p' DDD, o,p' DDE, p,p' DDE and percent moisture. As a quality control measure, the second jar from 14 randomly chosen sites was also analyzed. As a quality assurance measure, the second jar from five randomly chosen sites was analyzed by the Fish and Game Wildlife Water Pollution Control Laboratory of the California Department of Fish and Game. All results from the quality control and assurance samples showed good agreement with the analysis by the CDFA laboratory. The results of this survey are listed in Table III.

Examination of Table III shows several interesting features. The first is that contamination of agricultural soils in California by DDT_r is statewide. All sites sampled had DDT_r residues in the soil. The second feature is found on examination of the ratio of o,p' DDT + p,p' DDT to all DDT_r species. The ratio varies from 0 to 100 but is 49% on the average (see Figure 9). This means that on the average 49% of the detected DDT_r is present as DDT. This figure does not take into account the small amounts of DDD applied as a pesticide. These results are not unexpected. The values listed in scientific literature for the lifetime of DDT_r species in soil vary greatly but are usually 15 or more years.¹¹ The third feature is found when one examines the ratio of o,p' DDT to o,p' DDT + p,p' DDT. As stated earlier, this ratio was usually from 20 to 30% when the DDT was applied. Examination of Table I shows that, for most sites sampled, this ratio is still close to the value on application. The average value

for this ratio is 19% in this survey (see Figure 10). These levels of o,p' DDT are not unexpected for soils. The scientific literature shows that the o,p' isomer of DDT can be, and often is, longer lived than the p,p' isomer in the environment of soil.¹² The same scientific literature also shows that the o,p' isomer seems to be shorter-lived in the aquatic environment than the p,p' isomer. The argument has been made by some that a comparison of the ratios of the concentrations of various DDT isomers could be used to show how recently an application of DDT had been made. Clearly with both extremes occurring, o,p' and p,p' DDT long-lived in soil and short-lived in aquatic sediment, and with all the possible combinations of these extremes which can occur, such use of these ratios is tenuous at best. Additionally, since values of the o,p' isomer in soil of around 19% were the norm and not the exception, if illegal use of DDT is occurring it is on a statewide scale occurring everywhere at once. Such widespread illegal use could not occur without detection.

TABLE III
Data from 1985 Soil Monitoring survey for DDT

County Name	SiteN o.	% Mois- ture	o.p. DDT	p.p. DDT	o.p. DDD	p.p. DDD	o.p. DDE	p.p. DDE	o.p. DDI	p.p. DDI	o.p. DDE	p.p. DDE	Ratio ^b o.p./DDI (o.p.+ppp)/DDI	Ratio ^b DDI DDTr
Alameda	1	10	6	41	3	3	0	27					13	59
	2	9	202	862	92	210	24	471					19	57
Colusa	1	22	7	35	3	4	2	70					17	35
	2	32	2	9	1	2	3	38					18	20
Contra Costa	1	13	494	2,123	162	376	100	3,750					19	37
	2	1	81	431	9	97	24	465					16	46
Fresno	1	9	50	190	14	36	11	156					21	53
	2	11	20	50	3	7	6	94					29	39
	3	15	47	186	14	34	14	190					20	48
	4	9	195	717	24	NA	40	49					21	39
Glenn	1	22	32	228	25	NA	21	275					12	-
	2	10	15	54	4	10	2	93					22	39
Imperial	1	4	21	83	20	14	8	343					20	21
	2	4	12	32	17	15	5	165					27	18
	3	(4)d	-19	-36	-15	-17	-4	-182					-35	-20
	4	1	6	21	4	3	4	97					22	20
Kern	1	8	463	1,586	79	162	68	2,269					23	44
	2	-8	-463	-1,586	-79	-162	-68	-2,269					-23	-44
	3	6	37	40	0	4	2	51					48	57
	4	5	4	2	0	0	2	3					67	55
	4	6	7	52	0	4	10	76				12	40	

County Name	SiteN o.	% Mois- ture	o.p' DDT*	p.p' DDT*	o.p' DDD*	p.p' DDD*	o.p' DDE*	p.p' DDE*	Ratio ^b $\frac{o.p' DDI}{(o.p' + p.p') DDI}$	Ratio ^b $\frac{DDI}{DDTr}$
Kings	1	2	0	2	0	0	0	2	0	50
	2	13	15	35	2	11	9	89	30	31
Los Angeles	1	10	713	4,168	50	249	34	1,078	15	78
	2	4	2,326	13,740	459	629	654	9,553	14	59
	3	3	2,634	14,566	445	536	642	12,214	15	55
Madera	1	6	1	10	0	0	2	35	9	23
	2	8	0	1	0	0	0	0	0	100
	3	6	2	10	0	0	0	8	17	60
	4	8	26	224	3	25	3	59	10	74
		-8	-35	-255	-5	-27	-3	-69	-12	-74
Merced	1	5	99	535	18	53	24	1,082	16	35
	2	7	8	30	1	3	1	46	21	43
	3	8	3	12	0	0	0	18	20	45
	4	12	276	1,281	29	168	31	669	18	63
Monterey	1	16	508	1,285	85	186	37	1,116	28	56
	2	11	379	1,303	63	131	35	976	23	58
	3	10	262	926	41	199	16	717	22	55
	4	-10	-289	-1,059	-36	-146	-14	-719	-21	-60
		16	247	979	37	187	26	623	20	58
		-15	-282	-911	-40	-148	-16	-643	-24	-58
Napa	1	11	173	803	133	172	32	1,224	18	38
	2	10	233	1,607	185	179	26	1,046	13	56

County Name	Site No.	% Moisture	o.p' DDT	p.p' DDT	o.p' DDD	p.p' DDD	o.p' DDE	p.p' DDE	Ratio ^b o.p' DDI (o.p' + p.p')/DDT	Ratio ^b DDI DDTr
Orange	1	7	159	809	74	143	20	538	16	56
		-7	-150	-829	-76	-124	-18	-558	-15	-56
	2	7	23	149	9	25	3	112	13	54
		-6	-44	-161	-11	-37	-2	-114	-21	-56
	3	4	444	1,227	63	284	34	906	27	56
Placer	1	6	405	2,569	62	201	43	2,330	14	53
	2	18	14	129	0	7	0	11	10	89
Riverside	1	1	59	378	12	17	8	591	14	41
		-1	-52	-309	-11	-21	-7	-568	-14	-37
	2	4	63	163	35	20	24	1,219	28	15
	3	2	17	24	9	23	2	60	41	30
	4	0	28	81	7	7	12	311	26	24
Sacramento	1	10	19	83	2	12	1	67	19	55
	2	8	0	0	0	0	0	1	19	0
San Benito	1	23	49	177	7	19	13	214	22	47
	2	15	150	482	31	71	20	693	24	44
	3	16	64	574	52	107	6	618	10	45
	4	18	20	125	4	8	3	215	14	39
San Bernardino	1	1	3	10	0	0	0	15	23	45
	2	4	59	227	16	25	6	442	21	37
		-5	-59	-257	-9	-21	-7	-442	-19	-40
	3	12	6	44	19	32	6	158	12	19
		-4	-11	-49	-17	-40	-7	-221	-18	-18
	4	12	154	890	23	49	13	477	15	65
		-5	-142	-836	-10	-47	-14	-461	-15	-65

County Name	Site No.	% Moisture	o.p' DDT	P.p' DDT	o.p' DDD	P.p' DDD	o.p' DDE	P.p' DDE	Ratio ^b o.p' DDI (o.p' + p.p') DDI	Ratio ^b DDI DDT
San Diego	1	8	153	976	23	96	10	314	14	72
	2	4	7	31	2	4	1	33	18	49
	3	2	26	90	2	5	2	136	22	44
	4	-1	-24	-87	-5	-5	-2	-137	-22	-43
San Joaquin	1	4	50	511	26	97	6	289	9	57
	2	5	3	15	0	1	0	19	17	47
San Luis Obispo	1	5	72	322	13	41	7	258	18	55
	2	9	144	605	21	82	10	384	19	60
	3	21	606	2,661	124	370	52	1,688	19	59
	4	15	880	3,484	220	455	105	3,326	20	52
Santa Barbara	1	2	289	1,602	35	106	11	494	15	75
	2	2	444	2,121	47	179	11	627	17	75
	3	-2	-361	-1,706	-34	-133	-15	-516	-17	-75
	4	8	19	101	2	5	2	84	16	56
Santa Cruz	1	-6	-21	-116	-2	-10	-2	-94	-15	-56
	2	20	830	3,802	170	719	55	1,340	18	67
	3	18	910	3,438	96	411	44	1,446	21	69
	4	19	1,658	4,766	248	852	104	3,019	26	60
Solano	1	14	129	989	98	159	12	758	12	52
	2	9	22	93	5	12	4	168	19	38
Sonoma	1	11	1	6	0	0	0	4	14	64
	2	13	2	15	3	6	0	23	12	35
	1	7	154	853	563	247	31	516	15	43
	2	7	154	853	563	247	31	516	15	43

County Name	SiteNo	% Moisture	o,p' DDT	p,p' DDT	o,p' DDD	p,p' DDD	o,p' DDE	p,p' DDE	Ratio ^b o,p' DDT (o,p' + p,p')/DDT	Ratio ^b DDI DDT _T	
Stanislaus	1	5	26	79	5	NA	13	65	25	36	
	2	5	90	208	0	0	13	185	30	60	
	3	8	41	252	7	36	11	250	14	49	
	4	4	3	8	0	0	7	11	27	38	
Sutter	1	17	0	2	1	2	0	4	0	22	
	2	16	2	13	16	13	3	54	13	15	
Tehama	1	12	68	267	10	42	14	185	20	57	
	2	13	6	155	3	NA	8	61	4	-	
Tulare	1	6	3	33	1	3	1	48	8	40	
	2	7	14	132	6	15	1	104	10	54	
	3	9	17	84	6	10	3	197	17	32	
	4	12	27	131	7	7	18	35	17	70	
Ventura	1	4	321	1,347	40	160	29	828	19	61	
	2	2	182	466	30	99	10	379	28	56	
	3	-2	-184	-445	-27	-89	-9	-367	-29	-56	
	4	5	230	1,384	87	301	24	642	14	60	
Yolo	1	11	9	20	5	4	4	112	31	19	
	2	4	13	90	5	11	4	17	13	74	
	a. all values in parts per billion (dry weight)										
	b. all values times 100 (in percent)										
c. ratio of sum of MT isomers (o,p' + p,p') to sum of DDT + DDD + DOE isomers (o,p' + p,p') times 100 (in percent)											
d. values in parentheses are from analysis of duplicate samples collected at same site											




Figure 9. Percentage of o,p' and p,p' DDT in Total DDT Residues for All Counties Sampled.